

# Wnt signaling interacts with Shh to regulate taste papilla development

Ken Iwatsuki<sup>\*†</sup>, Hong-Xiang Liu<sup>‡</sup>, Albert Gründer<sup>§</sup>, Meredith A. Singer<sup>¶</sup>, Timothy F. Lane<sup>¶</sup>, Rudolf Grosschedl<sup>§</sup>, Charlotte M. Mistretta<sup>‡</sup>, and Robert F. Margolske<sup>\*†</sup>

<sup>\*</sup>Department of Neuroscience, Mount Sinai School of Medicine, 1425 Madison Avenue, New York, NY 10029; <sup>‡</sup>School of Dentistry, University of Michigan, Ann Arbor, MI 48109; <sup>§</sup>Max Planck Institute of Immunobiology, 79108 Freiburg, Germany; and <sup>¶</sup>Jonsson Comprehensive Cancer Center, University of California, Los Angeles, CA 90095

Edited by Jeremy Nathans, Johns Hopkins University School of Medicine, Baltimore, MD, and approved December 13, 2006 (received for review August 24, 2006)

Wnt and Shh signaling pathways are critical for the development and maturation of many epithelial tissues. Both pathways have roles in stem cell maintenance, tissue development, and tumorigenesis. However, linkage between these pathways in mammalian systems had not been well established. Here, we report that Shh expression in fungiform papillae and formation of normal mature fungiform papillae depend on signaling through Wnt and  $\beta$ -catenin. We observed that during fungiform papilla formation in mice, Shh and components of the Wnt/ $\beta$ -catenin signaling pathway are expressed together in the developing placode. The elimination of Wnt/ $\beta$ -catenin signaling in either Lef1 or Wnt10b knockout mice resulted in down-regulation of Shh expression. In addition, the size and number of fungiform papillae were greatly reduced in Lef1 knockout mice. By examining embryonic mouse tongues in culture we determined that activation of Wnt/ $\beta$ -catenin signaling up-regulates Shh expression. We observed that blocking Shh signaling in cultured tongue explants enhanced papillae formation and was accompanied by an up-regulation of Wnt/ $\beta$ -catenin signaling, indicating that Shh inhibits the Wnt/ $\beta$ -catenin pathway. Exogenously added Shh suppressed expression of endogenous Shh and inhibited Wnt/ $\beta$ -catenin signaling (assessed in TOPGAL mice), further implicating Shh as an inhibitor of the Wnt/ $\beta$ -catenin pathway. Our observations indicate that Wnt/ $\beta$ -catenin signaling and interactions between the Wnt and Shh pathways play essential roles in the development of fungiform papillae.

$\beta$ -catenin | Lef1 | Wnt10b | fungiform papilla

Wnt and Hedgehog (Hh) signaling molecules are indispensable for normal development of diverse epithelial tissues such as the gastrointestinal tract, hair follicles and teeth (for review, see refs. 1–3). Ectopic activation or elevated expression of these signaling molecules leads to tumorigenesis, whereas their inhibition causes developmental abnormalities in a variety of tissues (4, 5). Although both signaling pathways have been implicated often where epithelial–mesenchymal interactions take place (6, 7), demonstrations that these signaling pathways might interact or regulate each other are limited.

Taste tissue development in mice starts around embryonic day (E) 11.5, after the emergence of the tongue swelling on the floor of developing mandible (8, 9). This is followed by formation of the taste placode (E12.5), gustatory papillae (E13.5), and taste buds (around birth) (9). As with development of other epithelial tissues, formation and patterning of taste papillae are thought to be induced by epithelial–mesenchymal interactions (10). Expression of Shh, Bmp, Noggin, Fgf, and EGF is associated with taste papilla initiation and patterning, indicating the potential role of these factors in taste papilla development (11–14). Disruption of Shh signaling during the prenatal period by neutralizing antibody or by cyclopamine caused aberrant development and patterns of fungiform papillae (15, 16). Although Wnt signaling pathways are critical in the development of many epithelial tissues, they

had not been examined in the developing taste system, nor had their interaction with Shh been previously examined in the developing taste papillae.

## Results

**Expression of Wnt/ $\beta$ -Catenin Signaling Components in Developing Taste Organ.** To determine whether Wnt signaling is associated with taste papilla formation in mice, we monitored Wnt/ $\beta$ -catenin signaling activity and expression of Wnt signaling molecules and Wnt-activated transcriptional factors in developing taste tissue. We first examined Wnt/ $\beta$ -catenin signaling in TOPGAL mice that carry a  $\beta$ -galactosidase ( $\beta$ -gal) reporter gene under the control of an inducible promoter that responds to the complex of  $\beta$ -catenin with Tcf/Lef1 transcription factors (17). At E11.5, the developing tongue presents as paired lateral swellings with no obvious Wnt/ $\beta$ -catenin signaling apparent, as shown by an examination of  $\beta$ -gal activity in TOPGAL mice (Fig. 1 *A* and *F*).  $\beta$ -gal activity in TOPGAL mice could first be seen in the developing dorsal tongue at E12.5, the developmental stage when placode formation is initiated (Fig. 1 *B* and *G*). This  $\beta$ -gal activity was present in multiple sets of rows and was strongest in the row closest to the median sulcus (Fig. 1*B*).  $\beta$ -gal activity in the fungiform papillae is most intense at E14.5 then declines in overall intensity (Fig. 1 *C–E* and *H–J*) to become highly restricted to a small region of epithelium where taste buds develop (Fig. 1*J Inset*). At later stages,  $\beta$ -gal activity appeared in developing filiform papillae (Fig. 1 *D*, *E*, *I*, and *J*) and in the circumvallate (CV) papilla (noted by arrow heads in Fig. 1 *C–E Insets*).

We next used *in situ* hybridization to monitor expression of Wnt-regulated transcription factors between E12.5 and E14.5, the critical time for formation of the placode and subsequently fungiform papillae. Of the four Wnt-activated transcription factors (Tcf1, Tcf3, Tcf4, and Lef1) examined during these critical stages, only Lef1 showed an expression pattern similar to that of the Wnt/ $\beta$ -catenin signaling detected by  $\beta$ -gal activity in TOPGAL mice [compare Lef1 (Fig. 1 *L* and *M*) with  $\beta$ -gal (Fig.

Author contributions: K.I. and R.F.M. designed research; K.I., H.-X.L., and A.G. performed research; H.-X.L., A.G., M.A.S., T.F.L., R.G., and C.M.M. contributed new reagents/analytic tools; K.I., H.-X.L., C.M.M., and R.F.M. analyzed data; and K.I. and R.F.M. wrote the paper.

Conflict of interest statement: R.F.M. has a personal financial interest in the form of stock ownership in Linguagen Corp., receives consulting fees from Linguagen Corp., and is an inventor on patents and patent applications in the area of taste signal transduction that have been licensed to Linguagen Corp. Linguagen Corp. carries out work in the area of taste modification and taste signaling. The work described in the present manuscript would appear to be only peripherally related to the interests of Linguagen Corp.

This article is a PNAS direct submission.


Freely available online through the PNAS open access option.

Abbreviations: CV, circumvallate; En, embryonic day *n*.

<sup>†</sup>To whom correspondence should be addressed. E-mail: ken.iwatsuki@mssm.edu.


This article contains supporting information online at [www.pnas.org/cgi/content/full/0607399104/DC1](http://www.pnas.org/cgi/content/full/0607399104/DC1).

© 2007 by The National Academy of Sciences of the USA










**Fig. 3.** Ablating the Wnt/ $\beta$ -catenin pathway disrupts fungiform papillae development and reduces Shh expression. Fungiform papillae in the developing tongue were examined by SEM of E14 (A, B, M, and N) or postnatal day 0 (C and D) stage mice. The fungiform papillae at E14 of either Lef1 or Wnt10b null mice are reduced in size (B and N) in comparison to those of Lef1<sup>+/-</sup> heterozygous (A) or wild-type (M) mice. Lef1 null mice at postnatal day 0 show a marked decrease in the size and number of fungiform papillae (D; asterisks indicate missing or atrophied papillae, and arrows indicate remaining papillae), in comparison to those of the wild-type mice at postnatal day 0 (C; arrows mark fungiform papillae). Shh expression in developing tongues from wild-type and null mice was monitored by immunohistochemistry in whole mounts of tongues and mandibles (E–H and O–R) and in sections (I–L). Reduced Shh expression was observed in the fungiform papillae of developing tongues from Lef1 null mice (G, H, and L) and Wnt10b null mice (Q and R) compared with wild-type mice (E, F, J, O, and P). Normal Shh expression was observed in the CV papillae of Lef1 null (G and K, arrowheads), Wnt10b null (Q, arrowhead), and wild-type (E, I, and O, arrowheads) mice. (A–L) C57B/6 mice. (M–R) FVB mice. (Scale bars: A–D, M, and N, 200  $\mu$ m; E–H and O–R, 500  $\mu$ m; I–L, 100  $\mu$ m.)

**Wnt/ $\beta$ -Catenin Signaling Increases Fungiform Papillae Number.** To more closely examine the interactions of Wnt and Shh signaling molecules during taste papilla development, we turned to organ culture of embryonic tongues. Rodent tongues in culture have been shown to closely mimic *in vivo* spatial and temporal stages of morphogenesis and molecular expression, and to support the development of taste papillae (22, 23). Because the expression of Shh and Wnt signaling components in mice is seen only after E12.5, we maintained stage E12.5 (or E11.5; data not shown) tongues in culture for up to 2 days under varying treatment conditions. When we treated cultured E12.5 tongues with LiCl, a treatment that activates Wnt/ $\beta$ -catenin signaling through inhibition of glycogen synthase kinase-3 $\beta$  (GSK3 $\beta$ ) (24), we observed a dramatic increase in the number of fungiform



**Fig. 4.** Wnt and Shh interact in the developing tongue. Organ culture of E12.5 tongues from wild-type (*Left*) or TOPGAL (*Right*) mice treated for 2 days in culture with 5 mM LiCl (*A*), 50  $\mu$ g/ml anti-Shh antibody (5E1) (*C*), or 2.5  $\mu$ g/ml Shh-N (*E*) and examined for expression of Shh and  $\beta$ -gal in fungiform papillae. Control tongues (*A*, *C*, and *E*) were cultured for 2 days under standard culture conditions (STAND). Treatment with LiCl (*A*) or the anti-Shh antibody (*C*) up-regulated expression of both Shh and  $\beta$ -gal. Treatment with Shh-N (*E*) nearly abolished expression of Shh and down-regulated expression of  $\beta$ -gal. (*B*, *D*, and *F*) Histograms of the percent increase/decrease of the number of the Shh-immunoreactive fungiform papillae in treated vs. standard culture conditions, corresponding to images in *A*, *C*, and *E*, respectively ( $n = 6-7$ ; \*,  $P < 0.001$ ; means  $\pm$  SD). (Scale bar: 0.5 mm.)

papillae and number of Shh- and  $\beta$ -gal-expressing loci [compare tongues cultured under standard (STAND) vs. LiCl conditions] (Fig. 4 A and B). Control treatment with NaCl in place of LiCl did not alter Shh or  $\beta$ -gal expression (see SI Fig. 8), demonstrating the specificity of LiCl and indicating that it is activation of Wnt/ $\beta$ -catenin signaling by LiCl that increased both fungiform papillae number and their expression of Shh. Using *in situ* hybridization, we observed an increase in Shh mRNA after LiCl stimulation (data not shown), indicating that the up-regulation of Shh does not occur at the translational/posttranslational level. These results, in conjunction with the *in vivo* results from the Lef1 null mice, suggest that activation of Wnt/ $\beta$ -catenin signaling is necessary for both fungiform papillae formation and their expression of Shh.

**Inhibition of Shh Signaling Increases Wnt/ $\beta$ -Catenin Signaling and Fungiform Papillae Formation.** Previous reports with rodent tongue cultures showed that neutralizing Shh with a monoclonal anti-



body (5E1) or blocking Shh's activity with cyclopamine enhanced taste papillae formation (both size and number) accompanied by increased expression of Shh (15, 16, 18). Because we have seen coexpression of Wnt signaling elements and Shh, and because we determined that Wnt/ $\beta$ -catenin signaling up-regulates Shh expression, we set out to more closely examine interactions of these two signaling pathways. Blocking Shh function in TOPGAL mice with either the 5E1 anti-Shh antibody or with cyclopamine both elevated levels of Shh expression and enhanced expression of  $\beta$ -gal in fungiform papillae (Fig. 4 C and D and SI Fig. 9).

These results suggest that endogenous Shh may inhibit the Wnt/ $\beta$ -catenin signaling pathway. In turn, inhibition of Wnt by Shh would provide negative feedback regulation of Shh expression. As an initial test of this hypothesis, we treated cultured embryonic tongues with the purified N-terminal domain of mouse Shh (Shh-N), which is known to possess the full biological activity of Shh (25, 26), then examined its effects on Shh expression and Wnt/ $\beta$ -catenin signaling. We found that culturing E12.5 tongues for 2 days with Shh-N nearly abolished Shh expression in fungiform papillae (Fig. 4 E and F). By *in situ* hybridization we determined that this suppression of Shh was effected by decreased mRNA levels (SI Fig. 10). In addition, E12.5 tongues cultured in the presence of Shh-N had significantly reduced  $\beta$ -gal expression (i.e., Wnt/ $\beta$ -catenin signaling) (Fig. 4E). Together, these results suggest that Shh could directly or indirectly block fungiform papilla formation by inhibiting Wnt/ $\beta$ -catenin signaling.

To further test our hypothesis that endogenous Shh inhibits the Wnt/ $\beta$ -catenin signaling pathway during fungiform papillae development, we examined effects of modulating Shh expression in cultured tongues from Lef1 null embryos. In contrast to our results with wild-type mice, wherein the addition of Shh inhibitors markedly elevated Shh expression in fungiform papillae (Fig. 4 C and D), there was no such effect on embryonic tongues from Lef1 null mice in culture from the addition of 5E1 (SI Fig. 11) or cyclopamine (data not shown).

## Discussion

Wnts and Wnt signaling pathways are well known to be critical for the development of many epithelial tissues; however, they had not previously been known to play a role in the developing taste system. We initially detected Wnt/ $\beta$ -catenin signaling during fungiform papilla development by the use of TOPGAL reporter mice. We then confirmed expression of Wnt ligands in developing fungiform papillae by *in situ* hybridization and immunohistochemistry. Among the several Wnt ligands tested for expression in the developing taste organ, only Wnt10b was found to be selectively expressed in developing taste papillae where  $\beta$ -gal expression was detected in TOPGAL mice. Similarly, Lef1 was the only transcription factor among the Wnt-activated family of transcription factors to display such a pattern of expression. Thus, Lef1 and Wnt10b are the likely transcription factor and ligand, respectively, mediating Wnt/ $\beta$ -catenin signaling during fungiform papillae formation. Although Wnt10b has not been well characterized biochemically, there are several examples demonstrating Wnt10b's function as a ligand for the Wnt/ $\beta$ -catenin signaling pathway (27–30). In addition, we have confirmed that Wnt10b induces double axis formation when expressed in *Xenopus* embryos (K.I. and R.F.M., unpublished observations).

Shh is well documented to play an important role in the development of fungiform papillae (18). Yet, the underlying mechanism whereby Shh regulates taste papilla development was not known; nor was it known how disruption of Shh during fungiform papilla development led to increased expression of Shh. From our studies using knockout mice, we observed that ablation of Wnt/ $\beta$ -catenin signaling (by elimination of either




Fig. 5. Schematic diagram of Wnt–Shh regulatory loop during fungiform papillae formation.

Lef1 or Wnt10b) disrupted development of fungiform papillae and greatly reduced Shh expression therein. Also, up-regulation of the Wnt/ $\beta$ -catenin signaling by LiCl increased the number of fungiform papillae and their expression of Shh while addition of Shh suppressed Wnt/ $\beta$ -catenin signaling during fungiform papillae development in experiments using tongue cultures. These results indicate that interactions between Wnt and Shh signaling pathways play a role in fungiform papilla development.

There are precedents from developmental systems for interconnections between Wnt and Shh pathways. Experiments in *Drosophila* have shown that Wnt and Shh interactions occur in the embryonic epidermis and in the larval leg imaginal disk (31, 32). In the developing vertebrate spinal cord, interactions have been reported between Shh and Wnt signaling pathways in regulating patterning and cell specifications (32, 33). In mammalian systems, Lef1 null mice had been observed to have reduced expression of Shh in the developing tooth germ (34), and  $\beta$ -catenin mutant mice showed reduced Shh expression in hair follicles (35). In another example, Shh was shown to suppress Wnt10b mRNA expression in embryonic mandible cultures (36). Furthermore, it is intriguing that overexpression of active  $\beta$ -catenin causes ectopic expression of Shh during formation of hair follicles in mice and feathers in chickens (37–39).

We think it likely that Wnt/ $\beta$ -catenin signaling regulates Shh expression during development in fungiform papillae and in other epithelial tissues, and that in turn Shh suppresses the Wnt/ $\beta$ -catenin pathway in these tissues. At present we do not know whether Shh directly or indirectly suppresses the Wnt/ $\beta$ -catenin pathway, although we note that there are multiple Lef/Tcf binding sites within the upstream region of the mouse Shh gene (data not shown), suggesting that up-regulation of Shh mRNA may be a direct effect of activation of the Wnt/ $\beta$ -catenin pathway. We have developed a schematic model (Fig. 5) to describe the positive and negative feedback loops that may serve to coordinate Wnt and Shh pathways during normal development of fungiform papillae. In our model, Wnt/ $\beta$ -catenin signaling plays a central role in regulating Shh expression and the development of fungiform papillae, while Shh itself acts as an inhibitor of Wnt/ $\beta$ -catenin signaling.

In summary, our work indicates that Wnt and Shh signaling pathways are both essential for proper development of fungiform papillae, and that these two pathways interact via feedback circuits during the development of fungiform papillae.

## Materials and Methods

**Animals.** TOPGAL mice, originally generated by E. Fuchs (17), were purchased from The Jackson Laboratory and maintained in the CD-1 background. Lef1 null mice are described in ref. 20. Wnt10b<sup>−/−</sup> mice were created by T. F. Lane and P. Leder (29,

40). For timed pregnancies, the day that sperm were observed in the vaginal smear was designated 0.5 days postcoitum (E0.5).

**X-Gal Staining.** Tongues from TOPGAL mice at different developmental stages (E11.5, E12.5, E14.5, E16.5, and E18.5) were fixed in 0.5% glutaraldehyde in PBS for 10 min with rocking, washed in PBS, transferred into freshly prepared X-Gal solution (2.0 mM  $MgCl_2$ /0.01% sodium deoxycholate/0.02% Nonidet P-40/5 mM potassium ferricyanide/5 mM potassium ferrocyanide), and incubated at 37°C for 4 h. Stained tissues were photographed with a Zeiss dissecting microscope equipped with a Zeiss AxioCam HS digital camera system.

**Immunohistochemistry.** Antibodies used were rabbit anti-Lef1 antibody (20), goat anti-Shh-N (R & D Systems), anti- $\beta$ -catenin (BD Biosciences), and anti- $\beta$ -galactosidase (BioTrend). Tongues were fixed for 2 h at 4°C in 4% paraformaldehyde in PBS, transferred to 30% sucrose in PBS, and stored at 4°C overnight. Frozen sections (12  $\mu$ m) were rinsed in PBS and blocked in 0.3% Triton X-100 and 2% donkey serum in PBS for 1 h at room temperature. The primary antibodies were added to the section in a humidified chamber and incubated overnight at 4°C. Sections were washed in PBS, incubated with the secondary antibodies (Alexa Fluor 488 or 594 IgG against primary antibodies from Molecular Probes/Invitrogen; incubation continued for 60 min), and rinsed in PBS. For immunodetection in whole embryonic tongue, tongues in organ culture were fixed in 4% paraformaldehyde for 2 h and incubated in 6%  $H_2O_2$  for 2 h. After antigen retrieval with antigen unmasking solution (Vector Laboratories) for 10 min at 98°C, tongues were blocked with 0.3% Triton X-100 and 2% goat serum for 1 h at room temperature. Tongues were incubated overnight at 4°C with the antibodies against Shh-N (1  $\mu$ g/ml). After washing with 0.1% Triton X-100, tongues were treated with the VECTASTAIN ABC kit for goat (Vector Laboratories), and the Biotin-Streptavidin-HRP complex was detected by DAB substrate kit for peroxidase (Vector Laboratories). The peroxidase reaction was stopped by rinsing with PBS followed by 2% paraformaldehyde fixation. Stained tissues were photographed as described above. Numbers of Shh-immunoreactive fungiform papillae were analyzed statistically across the control and experimental

conditions by using the unpaired *t* test. A *P* value of <0.05 was considered significant.

**In Situ Hybridization.** Shh cDNA was from H. Kimura (St. Jude Children's Research Hospital, Memphis, TN). Wnt10b cDNA was from D. Agalliu (Columbia University, New York, NY). Lef1 cDNA was cloned by RT-PCR using cDNA from CV papillae. Tongue tissues frozen in OCT compound from E11.5, E12.5, E14.5, E16.5, and E18.5 stage mice were cut into sections (12  $\mu$ m) and subjected to postfixation (10 min in 4% paraformaldehyde, followed by acetylation in acetic anhydride for 10 min). After three washes in PBS, sections were prehybridized in hybridization buffer (5 $\times$  SSC/50% formamide/1 $\times$  Denhardt's solution/1 mg/ml salmon sperm DNA/1 mg/ml tRNA). Hybridizations were performed with digoxigenin-labeled cRNA probes in the hybridization buffer for 18 h at 72°C. Hybridization signals were detected by alkaline phosphatase-conjugated anti-digoxigenin antibodies plus NBT/BCIP substrate (Roche).

**SEM Analysis.** SEM was performed as described in ref. 15.

**Organ Culture of the Tongue.** Organ culture of tongues from rodent embryos was described in ref. 15. Embryonic tongues from E11.5 or E12.5 stage mice were cultured on Millipore Millicell-HA culture plate inserts in standard (STAND) culture medium [DMEM/F12 medium supplemented with 1% FBS, 2% B27 culture supplement (GIBCO/Invitrogen), and 50  $\mu$ g/ml gentamicin sulfate] for 2 days. To activate Wnt/ $\beta$ -catenin signaling, 5 mM LiCl was supplemented to the STAND medium. To disrupt Shh signaling, either Shh-blocking antibody 5E1 (50  $\mu$ g/ml; Developmental Studies Hybridoma Bank) or cyclopamine (5  $\mu$ M; Toronto Research Chemicals) was added to the STAND medium. To induce Shh signaling, 2.5  $\mu$ g/ml Shh N-terminal peptide (Shh-N; R & D Systems) was supplemented to the STAND medium.

We thank Dr. H. Kimura for Shh cDNA, Dr. D. Agalliu for Wnt10b cDNA, and N. Harekaki and R. Takubo for technical support. This work was supported in part by National Institutes of Health Grants DC003055 (to R.F.M.), DC003155 (to R.F.M.), and DC00456 (to C.M.M.) and a fellowship from the Japan Society for the Promotion of Science (to K.I.).

- Hooper JE, Scott MP (2005) *Nat Rev Mol Cell Biol* 6:306–317.
- Logan CY, Nusse R (2004) *Annu Rev Cell Dev Biol* 20:781–810.
- Gregorieff A, Clevers H (2005) *Genes Dev* 19:877–890.
- Taipale J, Beachy PA (2001) *Nature* 411:349–354.
- McMahon AP, Ingham PW, Tabin CJ (2003) *Curr Top Dev Biol* 53:1–114.
- Bitgood MJ, McMahon AP (1995) *Dev Biol* 172:126–138.
- Reddy S, Andl T, Bagasra A, Lu MM, Epstein DJ, Morrissy EE, Millar SE (2001) *Mech Dev* 107:69–82.
- Paulson RB, Hayes TG, Sucheston ME (1985) *J Craniofac Genet Dev Biol* 5:59–73.
- Kaufman MH (1992) *The Atlas of Mouse Development* (Academic, London).
- Mistretta CM (1998) *Ann NY Acad Sci* 855:1–13.
- Jung HS, Oropeza V, Thesleff I (1999) *Mech Dev* 81:179–182.
- Hall JM, Hooper JE, Finger TE (1999) *J Comp Neurol* 406:143–155.
- Zhou Y, Liu HX, Mistretta CM (2006) *Dev Biol* 297:198–213.
- Mistretta CM, Grigaliunas A, Liu HX (2005) *Chem Senses* 30(Suppl 1):i52–i53.
- Mistretta CM, Liu HX, Gaffield W, MacCallum DK (2003) *Dev Biol* 254:1–18.
- Hall JM, Bell ML, Finger TE (2003) *Dev Biol* 255:263–277.
- DasGupta R, Fuchs E (1999) *Development (Cambridge, UK)* 126:4557–4568.
- Liu HX, MacCallum DK, Edwards C, Gaffield W, Mistretta CM (2004) *Dev Biol* 276:280–300.
- Oosterwegel M, van de Wetering M, Timmerman J, Kruisbeek A, Destree O, Meijlink F, Clevers H (1993) *Development (Cambridge, UK)* 118:439–448.
- van Genderen C, Okamura RM, Farinas I, Quo RG, Parslow TG, Bruhn L, Grosschedl R (1994) *Genes Dev* 8:2691–2703.
- Zhou P, Byrne C, Jacobs J, Fuchs E (1995) *Genes Dev* 9:700–713.
- Mbiene JP, MacCallum DK, Mistretta CM (1997) *J Comp Neurol* 377:324–340.
- Nosrat CA, MacCallum DK, Mistretta CM (2001) *Cell Tissue Res* 303:35–45.
- Klein PS, Melton DA (1996) *Proc Natl Acad Sci USA* 93:8455–8459.
- Lee JJ, Ekker SC, von Kessler DP, Porter JA, Sun BI, Beachy PA (1994) *Science* 266:1528–1537.
- Ingham PW, McMahon AP (2001) *Genes Dev* 15:3059–3087.
- Lane TF, Leder P (1997) *Oncogene* 15:2133–2144.
- Ross SE, Hemati N, Longo KA, Bennett CN, Lucas PC, Erickson RL, MacDougald OA (2000) *Science* 289:950–953.
- Bennett CN, Longo KA, Wright WS, Suva LJ, Lane TF, Hankenson KD, MacDougald OA (2005) *Proc Natl Acad Sci USA* 102:3324–3329.
- Ouji Y, Yoshikawa M, Shiroy A, Ishizaka S (2006) *Biochem Biophys Res Commun* 342:1063–1069.
- Heemskerk J, DiNardo S (1994) *Cell* 76:449–460.
- Roelink H (1996) *Curr Opin Neurobiol* 6:33–40.
- Lei Q, Jeong Y, Misra K, Li S, Zelman AK, Epstein DJ, Matisse MP (2006) *Dev Cell* 11:325–337.
- Kratochwil K, Dull M, Farinas I, Galceran J, Grosschedl R (1996) *Genes Dev* 10:1382–1394.
- Huelsken J, Vogel R, Erdmann B, Cotsarelis G, Birchmeier W (2001) *Cell* 105:533–545.
- Dassule HR, McMahon AP (1998) *Dev Biol* 202:215–227.
- Gat U, DasGupta R, Degenstein L, Fuchs E (1998) *Cell* 95:605–614.
- Silva-Vargas V, Lo Celso C, Giangreco A, Ofstad T, Prowse DM, Braun KM, Watt FM (2005) *Dev Cell* 9:121–131.
- Noramly S, Freeman A, Morgan BA (1999) *Development (Cambridge, UK)* 126:3509–3521.
- Vertino AM, Taylor-Jones JM, Longo KA, Bearden ED, Lane TF, McGehee RE, Jr, MacDougald OA, Peterson CA (2005) *Mol Biol Cell* 16:2039–2048.